
High-throughput screening assay for sickle cell disease drug discovery

A high-throughput screening assay for discovering small molecules that directly inhibit sickle hemoglobin oligomerization.



IP Status: US Patent Pending; Application No. 17/953,981

Applications

- Sickle cell drug discovery
- Sickle cell disease research

Key Benefits & Differentiators

- High-sensitivity: Assay uses high-sensitivity fluorescence lifetime (FLT) FRET
- High-throughput: Assay uses FLT plate reader technology enabling high-throughput screening

Technology Overview

Sickle cell disease (SCD) is a devastating hereditary disorder caused by a defect in the hemoglobin gene that leads to sickle hemoglobin (HbS) polymers that distort the shape of red blood cells. Despite contributing to a shortened lifespan, high infant mortality rates, and significantly reduced quality of life, treatment options for SCD patients are limited to four FDA-approved therapies that do not dramatically improve outcomes. While gene therapy has elicited considerable excitement, the need for bone marrow ablation and the high cost preclude broad application of this approach. Millions of SCD patients worldwide would benefit from new, affordable, and easily accessible therapies. Thus, new small molecule approaches are desperately needed, but the pipeline for identifying and validating new therapies is extremely limited.

After observing that deoxygenated HbS forms temporally stable oligomers at sub-nucleation concentrations, researchers at the University of Minnesota developed a high-throughput

Technology ID 2019-329

Category

Engineering & Physical Sciences/Processes Life Sciences/Biochemicals & Small Molecules Life Sciences/Human Health Life Sciences/Pharmaceuticals Life Sciences/Research Tools Life Sciences/Therapeutics

View online page



screening (HTS) assay for discovering small molecules that directly inhibit HbS oligomerization. This FRET-based assay leverages high-sensitivity fluorescence lifetime measurements that monitor the temporally stable HbS oligomers. In the lab, the assay demonstrated HbS oligomer sensitivity to known polymerization inhibitors and identified novel HbS oligomerization inhibitors that reduced hypoxia-induced impairment of sickle blood flow without increasing hemoglobin affinity. This screening platform has commercial potential for large-scale HTS of small molecules that could be developed into new therapeutics for SCD.

Phase of Development

TRL: 4-5

High-throughput screening assay demonstrated in the lab HbS oligomer sensitivity to known polymerization inhibitors and identified novel HbS oligomerization inhibitors.

Desired Partnerships

This technology is now available for:

- License
- Sponsored research
- Co-development

Please contact our office to share your business' needs and learn more.

Researchers

- David Wood, PhD Associate Professor, Department of Biomedical Engineering
- Jonathan Sachs, PhD Professor, Department of Biomedical Engineering

References

 Nagamani VunnaNagamani Vunnam , Scott Hansen, Dillon C. Williams, MaryJane Olivia Been, Chih Hung Lo, Anil K. Pandey, Carolyn N. Paulson, John A. Rohde, David D. Thomas, Jonathan N. Sachs, and David K. Wood(2022) , https://doi.org/10.1021/acs.biomac.2c00671, https://pubs.acs.org/doi/10.1021/acs.biomac.2c00671, 23, 9, 3822-3830